

EFFECTIVE
2/6/08

NEBRASKA DEPARTMENT OF HEALTH
AND HUMAN SERVICES

173 NAC 4
Attachment 1

CHILDHOOD IMMUNIZATION SCHEDULE

Child's Age At Entrance or At Time of Record Review	<<<<Minimum>>>> Number of Doses Required of Each Vaccine						
	DTaP/DT/ Td/Tdap Vaccine	Polio Vaccine	MMR/ MMRV Vaccine	Hib Vaccine	Hep B Vaccine	VZV Vaccine	PCV Vaccine
0 but not yet 3 Months	None	None	None	None	None	None	None
3 months but not yet 5 Months	1 Dose	1 Dose	None	1 Dose	1 Dose	None	1 Dose
5 months but not yet 7 Months	2 Doses	2 Doses	None	2 Doses	2 Doses	None	2 Doses
7 months but not yet 16 Months	3 Doses	2 Doses	None	2 Doses **	3 Doses	None	3 Doses ****
16 months but not yet 19 Months	3 Doses	2 Doses	1 Dose *	3 Doses **	3 Doses	1 Dose or Document ed History of Disease	4 Doses ****
19 Months to School Entry	4 Doses	3 Doses	1 Dose *	3 Doses **	3 Doses	1 Dose or Document ed History of Disease	4 Doses ****
At School Entry	4 Doses	3 Doses	2 Doses *	None ***	3 Doses	2 Doses or Document ed History of Disease	None

* First dose must be given no earlier than 4 days before the 1st birthday.

** Hib vaccine is recommended to be given in a multiple dose schedule beginning in infancy. However, any child who has received a single dose of Hib vaccine at or after 15 months of age is appropriately immunized.

*** Hib vaccine is not required after child reaches 5 years of age.

**** If a child is 7 months of age or older when they start the PCV series, they do not need all 4 doses of PCV. Unvaccinated children who begin the PCV series between 7-11 months of age should receive 2 doses 4 weeks apart with a booster at 12-15 months. Unvaccinated children who begin the PCV series between 12-23 months of age should receive 2 doses 8 weeks apart. Unvaccinated children 24-59 months of age need 1 dose of PCV. Children with high-risk conditions age 24-59 months should follow their physician's advice.